




Motivation and background



Andersen et al (2017) present a mechanism-based mathematical model describing the hematopoietic (blood-producing) system of the human body during the development of the Myeloproliferative Neoplasms (MPNs), a group of slowly developing blood cancers. The model consists of a six-dimensional system of ODEs.

Through time-scale considerations, the model can be reduced to a simpler two-dimensional system describing the slow dynamics (Ottesen et al, 2019).


So far, model analysis has granted some insight about the cytokines related to MPNs as well as prediction of patient-specific treatment response.

However, some changes to the model are currently being considered:

- A subset of patients display response that are not readily reproducible by the present model formulation.
- The treatment considered (INF- α) is known to affect features of the hematopoietic system not currently included in the model.


Mechanisms of stem cells

- The figure depicts a compartment diagram of a separate model considering solely the hematopoietic stem cells within the bone marrow micro-environment.
- Inspired by the work of Wang et al (2017), the model considers competition between stem cells in a finite stem cell niche of capacity K .
- Stem cells bound to the niche are considered quiescent while free stem cells can divide, differentiate or return to quiescence by rebinding with the niche.



N : Niche-bound; A : Free, active; I : Free, inactive. C and L denote cell-types.

Results



- Time-scale considerations allow for a reduction of the stem cell model to a system of two ODEs, modelling the slow time-scale.

$$\dot{N}_C = \left(\frac{2(1 - N_C - N_L)}{\alpha_C + 1 - N_C - N_L} \left(\frac{r_C}{r_C + d_{CA}} \right) - 1 \right) N_C$$

$$\dot{N}_L = \mu \left(\frac{2(1 - N_C - N_L)}{\alpha_L + 1 - N_C - N_L} \left(\frac{r_L}{r_L + d_{LA}} \right) - 1 \right) N_L$$

Where $\alpha_C = d_{CI}/(b_{CI}K)$, $\alpha_L = d_{LI}/(b_{LI}K)$ and $\mu = u_L/u_C$

- Stem cell compartments in the Cancitis model can be replaced with the slow time-scale stem cell model leading to an extended model.
- The figure displays disease development, two examples of data for patients undergoing treatment as well as simulate treatment in the models.
- Both the normal and extended model can reproduce the responses to treatment, by step-wise changing parameters at treatment onset.

Preliminary conclusions

- When modelling treatment-response of MPN patients, the Cancitis model implies **increased elimination** of malignant cells as response to treatment.
- The model can be extended to include the **stem cell dynamics** more accurately, which allows for **alternative hypotheses about which parameters are affected by treatment**.
- Preliminary results suggests that **increased activation of quiescent stem cells** ($u_C \uparrow$ and $u_L \uparrow$) and increased differentiation ($d_{LI} \uparrow$) lead to the delayed responses seen in data.

References

- Andersen, M., Sajid, Z., Pedersen, R. K., Gudmand-Hoeyer, J., Ellervik, C., Skov, V., Kjaer, L., Pallisgaard, N., Kruse, T. A., Thomassen, M., Troelsen, J., Hasselbalch, H. C., Ottesen, J. T. (2017). Mathematical modelling as a proof of concept for MPNs as a human inflammation model for cancer development. PLOS ONE, 12(8), e0183620. <https://doi.org/10.1371/journal.pone.0183620>
- Ottesen, J. T., Pedersen, R. K., Sajid, Z., Gudmand-Hoeyer, J., Bangsgaard, K. O., Skov, V., Kjaer, L., Knudsen, T.A., Pallisgaard, N., Hasselbalch, H. C., Andersen, M. (2019). Bridging blood cancers and inflammation: The reduced hematopoietic stem cell frequency predicts outcome in acute myeloid leukemia. Haematologica, 102(9), 1567-1577. <https://doi.org/10.3324/haematol.2016.163584>